

AD-A150 555

ADSORPTION SITE ADSORPTION ENERGY AND NORMAL VIBRATION
FREQUENCY OF H₂NIC... (U) CORNELL UNIV ITHACA NY LAB OF
ATOMIC AND SOLID STATE PHYSICS.. C UMRIGAR ET AL.

1/1

UNCLASSIFIED

07 FEB 85 8 N00014-82-K-0576

F/G 7/4


NL



END

FORMED

DTN



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

4

AD-A150 555

DTIC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 8	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Adsorption Site, Adsorption Energy and Normal Vibration Frequency of H/Ni(100) via Total Energy Calculations		5. TYPE OF REPORT & PERIOD COVERED Interim
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Cyrus Umrigar and John W. Wilkins		8. CONTRACT OR GRANT NUMBER(s) N00014-82-K-0576
9. PERFORMING ORGANIZATION NAME AND ADDRESS Laboratory of Atomic and Solid State Physics Cornell University Ithaca, N.Y. 14853		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Washington, D. C. 10360		12. REPORT DATE February 7, 1985
		13. NUMBER OF PAGES 11
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Submitted to Physical Review Letters.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Density-functional total energy calculations of H/Ni(100) show that the equilibrium H position is in the center site, 0.6a ₀ above the plane of the surface Ni atoms (with a Ni-H bond length of 1.8 Å). The bridge-site minimum lies only 0.1 eV higher than the center site minimum, whereas the top site minimum lies about 0.3 eV higher. This is consistent with a high hydrogen surface mobility.		

**DTIC
ELECTE**
FEB 19 1985
D

DD FORM 1473

EDITION OF 1 NOV 55 IS OBSOLETE
S/N 0102-014-66011

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

0

Adsorption Site, Adsorption Energy and Normal Vibration Frequency of H/Ni(100) via Total Energy Calculations

Cyrus Umrigar and John. W. Wilkins

Laboratory of Atomic and Solid State Physics
Cornell University, Ithaca, NY 14853

ABSTRACT

Density-functional total energy calculations of H/Ni(100) show that the equilibrium H position is in the center site, 0.6 Å above the plane of the surface Ni atoms (with a Ni-H bond length of 1.8 Å). The bridge-site minimum lies only 0.1 eV higher than the center site minimum, whereas the top site minimum lies about 0.3 eV higher. This is consistent with a high hydrogen surface mobility

Accession For	
TIS GRA&I	<input checked="" type="checkbox"/>
TIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
Availability Codes	
Dist	Avail and/or Special
A-1	

In recent years it has become possible to use density-functional total energy calculations to predict a variety of properties of materials. Calculations have been performed to predict the lattice constant, cohesive energy and bulk moduli of solids^{1, 2}; high pressure phases of materials^{3, 4}; geometries, binding energies and vibration frequencies of molecules⁵; surface energies⁶; and the geometry of monolayer films⁷ and of surfaces.^{8, 9} Most total energy calculations of surfaces have been confined to clean semi-conductor surfaces, the reason being that the pseudo-potential method used in these calculations is better suited to semi-conductors than to transition metals, either clean or adsorbate covered. Very recently there has been some work on clean transition metal surfaces using all-electron methods.^{6, 9}



In this letter we present total energy calculations for an adsorbate on a transition metal surface. The particular system we have chosen is P(1x1) H/Ni(100) because a variety of experiments that we can compare with: thermal desorption¹⁰, electron energy loss spectroscopy¹¹, ion-channeling¹² and He scattering¹³ have been performed on this system. We have performed calculations to determine the geometry of the adsorbed H, its binding energy, its vibration frequency and the barrier to surface diffusion.

Our calculations were performed using the Linearized Augmented Plane Wave (LAPW) method.^{14, 15} The core electrons (Ni 1s-3p) were treated fully relativistically and the valence electrons scalar relativistically. The core is included in the calculation of the total energy. The potential was allowed to have a general form everywhere (no muffin-tin approximation), except that non-spherical terms within the muffin-tin spheres were evaluated approximately. The non-spin-polarized form of the Vosko-Wilk-Nusair (VWN)¹⁶ exchange-correlation potential was used. Some of the calculations were redone using the Hedin-Lundqvist (HL)¹⁷ potential in order to check the sensitivity of the results to the choice of the exchange-correlation potential. Most of the calculations were performed on 7 layer slabs consisting of 5 layers of Ni with a layer of H on each side.

Fig. 1 shows the adsorption energy $E_{ads} = (E_{H/Ni}^{slab} - E_{Ni}^{slab} - 2E_H^{atom})/2$ as a function of the vertical distance of the H atoms above the surface Ni plane for the center(C), bridge(B), and atop(A) sites. Here $E_{H/Ni}^{slab}$ and E_{Ni}^{slab} are, respectively, the total energies (per unit cell) of the H covered, and clean Ni slabs, and E_H^{atom} is the energy of a free spin-polarized H atom (0.957 Ry or 13.02 eV for VWN, 0.976 Ry or 13.28 eV for HL). The equilibrium position is found to be 0.6 a_o ($a_o = 0.529 \text{ \AA}$) above the center site (corresponding to a Ni-H bond length of 1.79 \AA). This is probably a small underestimate ($\approx 2\%$ or 0.07 a_o in the bond length, or 0.31 a_o in the vertical distance z) since local density functional

calculations for bulk Ni gives a bond length that is about 2% too small. Hence, we expect the true position of the H atoms to be at about $0.9 a_o$ above the surface Ni plane. Also the inclusion of magnetism (which we have neglected) would tend to increase interatomic distances slightly. In fact, a recent spin-polarized calculation¹⁸ gives $z = 0.69 a_o$. It is difficult to determine the position of H on surfaces experimentally,¹⁹ but there is in fact a reliable determination of the position of H on Ni(100) from transmission ion-channeling experiments.¹² These experiments show that H adsorbs $0.95 \pm 0.2 a_o$ above the center site, in excellent agreement with our estimate ($0.6 + 0.3 a_o$).

The calculated value of the adsorption energy (3.61 eV for VWN, 3.42 eV for HL) is considerably larger than the experimental value¹⁰ (2.7 eV). It appears to be a common failing of local density functional theory that it gives too much binding^{1, 5, 20} the reason being that the large underestimate of the total energies of the free atom and of the solid (4.3% for H, 0.1% for Ni) due to the local density approximation do not quite cancel, the error being slightly larger for the atom than for the (more homogeneous) solid.

The bridge site minimum lies only 0.08 eV higher than the center site minimum and the top site minimum lies about 0.32 eV higher. It ought to be mentioned here that although the individual curves have a relative numerical precision of better than 0.01 eV, there may be a relative error between the energies of the center and bridge site curves of 0.1 eV and of the center and top site curves of 0.2 eV. The reason for this is that since H atoms approach the surface Ni atoms more closely at the bridge and top sites, one is forced to choose smaller muffin-tin radii when calculating these curves, resulting in a less well converged basis set and greater uncertainties in the extrapolation to the infinite basis size energies. The extrapolation procedure is explained in greater detail later.

The near degeneracy of the center and bridge site minima implies that the H atoms will have large rms displacements from their equilibrium positions or may even be mobile on the surface. There is some experimental evidence that the barrier to surface diffusion of the H is small. Ion-channeling experiments¹² show that the H atoms have a rms displacement of $0.4a_0$ from their equilibrium position. Laser-induced desorption experiments²¹ show that the barrier to diffusion is about 0.17 eV for H on a laser treated Ni(100) surface. Also H has been observed to be mobile on other transition metal surfaces by field-ion microscopy.²²

Also shown in Fig. 1 are other theoretical results obtained using the generalized valence bond (GVB)²³ and the effective medium²⁴ methods. The inclusion of configuration effects in GVB restricts its application to small clusters of atoms which may not adequately represent an extended surface and may not represent the neighborhood of the center, bridge and top sites equally well. Also, the d electrons are not included explicitly. The effective medium technique is a simple method based on density functional theory. It has the advantage that it is fast enough to use in complicated geometries where the more accurate methods used in our work are not feasible and the disadvantage that it is approximate, in particular the interaction of the adsorbate with the metal d-electrons is treated in a crude way. The GVB calculation gives an equilibrium position in good agreement with ours but the relative energy difference of the minima of the curves for the three sites is much larger than in our calculation. The effective medium approach gives basically the same energy differences between the minima of the three curves as our calculation, and bond lengths that are all somewhat larger than ours. Also the detailed shape of the curves is somewhat different from ours. The semi-empirical embedded atom method²⁵ gives energy differences in good agreement with, and bond lengths in fair agreement with our results.

Electron energy loss (EELS) experiments¹¹ for a full layer of H on Ni(100) yield a vibration frequency of 78 meV. A precise calculation of this frequency would entail a calculation of the total energy at all points in space in the neighbourhood of the minimum energy surface. Then the difference of the energies of the two lowest eigenstates for the motion of the H atom as a whole would give the EELS vibration frequency. This has been done within the effective medium approach²⁶ but is difficult to do in our approach. However an approximate estimate can be obtained from the second derivative of the energy curve near the minimum. This gives a value of 90 meV which compares reasonably well with experiment given the approximations we have made.

Since the energies of interest are obtained by subtracting total energies which are several orders of magnitude larger than the energies of interest, it is important to test the convergence of our results as a function of the size of the various expansion sets used, and to test the sensitivity of our results to the choice of various parameters. Here, we discuss only some of the tests performed, namely convergence with respect to i) the size of the basis set used for expanding the wavefunctions, and ii) the number of layers in the film, and sensitivity to the choice of the exchange-correlation potential.

One virtue of the LAPW method is that the basis can be systematically improved by including all plane waves with wavevectors lying within successively larger spheres of radius K_{\max} . Fig. 2a shows the total energy versus the distance of the center site H atoms above surface Ni layer for two values of $R_{MT} K_{\max}$ where R_{MT} is the muffin tin radius of the center layer Ni atoms. In Fig. 2b the total energy for $z=0.6a_0$ is plotted versus $\exp(-R_{MT} K_{\max})$. The six points shown fit the exponential with an rms error of 0.008 eV. An almost equally good fit can be obtained to a power law (rms error 0.011 eV) with a exponent ≈ -12.5 . The estimate of E_{tot} for a fully converged basis (plotted as the bottom curve in

Fig. 2a) is obtained assuming the exponential dependence. Assuming power law dependence would give a curve uniformly shifted down by 0.08 eV. The position of the minimum and the shape of the curves for $R_{MT} K_{\max} = 7.5$ and 8.0 are almost identical. Hence it is possible to obtain the correct vertical position without having a fully converged basis as has been noticed by other authors.⁷ However, since the center, bridge, and top site curves are calculated with different values of R_{MT} and are therefore not equally well converged, it is very important to perform the extrapolation to obtain the relative energy differences between them.

The binding energies of H at two positions above the center site were recalculated using a thinner slab consisting of 3 Ni layers with a layer of H on each side (shown as square points in Fig. 1). The close agreement between these energies and those for the fatter slab give us confidence that the energies are converged with respect to the thickness of the slab.²⁷

Fig. 1 also shows that using the HL exchange-correlation potential instead of the VWN potential results in a binding energy curve almost parallel to that for the VWN potential over the range of distances shown but shifted up by 0.19 eV.

We have shown that accurate total energy calculations can be used to determine the geometry of adsorbates on surfaces.

It is a pleasure to acknowledge many useful discussions with Anders Carlsson, Gayanath Fernando, Matti Manninen, Jens Norskov, Mats Persson, and Mark Stiles and assistance with computing by the Center for Materials Science of Los Alamos National Laboratory. This research was funded by the Office of Naval Research.

References


1. V. L. Moruzzi, A. J. F. Janak, and A. R. Williams, Calculated Electronic Properties of Metals (Pergammon, 1978).
2. H. J. F. Jansen, and A. J. Freeman, Phys. Rev. B **30**, 561 (1984).
3. M. T. Yin, and M. L. Cohen, Phys. Rev. Lett. **50**, 200 (1983).
4. A. E. Carlsson, and J. W. Wilkins, Phys. Rev. B **29**, 5836 (1984).
5. J. E. Muller, J. Harris, and R.D. Jones, J. Chem. Phys. **79**, 1874 (1983).
6. R. Richter, J. Gay, and J. Smith, Proceedings of the First International Conference on the Structure of Surfaces 1984.
7. M. Weinert, E. Wimmer, and A. J. Freeman Phys. Rev. B **26**, 4571 (1982).
8. J. E. Northrup, and M. L. Cohen, Phys. Rev. B **29**, 5944 (1984).
9. C. L. Fu, S. Ohnishi, E. Wimmer, and A. J. Freeman Phys. Rev. Lett. **53**, 675 (1984).
10. K. Christman, O. Schober, G. Ertl, and M. Neumann J. Chem. Phys. **60**, 4528 (1974).
11. Experiments with a full layer of H yield a vibration frequency of 78 meV, S. Andersson, personal communication. Earlier experiments with a lower coverage gave a frequency of 74 meV, S. Andersson, Chem. Phys. Lett. **555**, 185 (1978).
12. I. Stensgaard and Jakobsen, to be published, and personal communication.
13. K. H. Rieder, and H. Wilsch, Surf. Sci. **131**, 245 (1983).
14. O. K. Andersen, Phys. Rev. B **12**, 3060 (1975).
15. M. Posternak, H. Krakauer, A.J. Freeman, D. D. Koelling, Phys. Rev. B **21**, 5601 (1980).


16. S. H. Vosko, L. Wilk, and M. Nusair, *Can. J. Phys.* **58**, 1200 (1980).
17. L. Hedin, and B. J. Lundqvist, *J. Phys. C* **4**, 2064 (1971).
18. M. Weinert and J.W. Davenport, *Bull. Am. Phys. Soc.* **29**, 434 (1984), and personal communication.
19. LEED is not a suitable technique because of the small cross-section of H for electrons. He scattering experiments have been performed and interpreted to give $z = 1.8 \pm 0.1 a_0$ (see Ref. 13) but we have doubts about the interpretation of the experiments.
20. It is pointed out in Ref. 7 that use of a limited basis gave a binding energy of a graphite monolayer in good agreement with experiment whereas a well converged basis gives about 1.3 eV/atom too much binding.
21. R. Hall, personal communication.
22. R. DiFoggio, and R. Gomer, *Phys. Rev. B* **25**, 3490 (1982).
23. Thomas H. Upton, and William A. Goddard, *J CRC Crit Rev in Sol. St and Mat. Sci.* **10**, 261 (1981).
24. J. K. Norskov, *Phys. Rev. Lett.* **48**, 1620 (1982), and personal communication.
25. M. S. Daw, and M. I. Baskes, *Phys. Rev. B* **29**, 6443 (1984).
26. M. J. Puska et. al., *Phys. Rev. Lett.* **51**, 1081 (1983).
27. A by-product of the clean Ni calculations is the determination of the value of the Ni(100) surface energy (1.0 eV/atom or $2.6 J/m^2$) deduced from the 3- and 5-layer total energies.

Figure Captions

Figure 1. Binding energy calculated by various authors as a function of the vertical distance of the H atoms above the surface Ni layer, in the center (—) bridge (— — —) and atop (...) sites. The calculated points are generally indicated by dots. The two squares show that the results for VWN 5-layer are similar to VWN 7-layer.

Figure 2. Convergence of the total energy E_{tot} with respect to the basis size. Fig. (2a) shows curves of E_{tot} as a function of the vertical distance of the H atoms above the surface Ni layer for $R_{MT} K_{max}$ equal to 7.5 and 8. R_{MT} is the muffin-tin radius of the central layer Ni atoms and K_{max} is the largest wavevector included in the basis functions. The estimated curve for an infinite basis is also shown. Fig.(2b) shows the procedure used to estimate E_{tot} for an infinite basis. $\alpha_{fit} = 1.78$





END

FILMED

3-85

DTIC